

[Click Here](#)



Drug interactions are reported among people who take Amoxicillin and clavulanate potassium (amoxicillin; clavulanate potassium) and Tylenol (acetaminophen). Common drug interactions include nausea among females and dizziness among males. The phase IV clinical study analyzes what interactions people have when they take Amoxicillin and clavulanate potassium and Tylenol, and groups them by gender, age and more. It is created by eHealthMe based on reports of 141 people who take the same drugs from the FDA, and is updated regularly. With medical big data and AI/ML algorithms, eHealthMe is running millions of data-driven phase IV clinical trials continuously. Our original studies have been referenced on 800+ peer-reviewed medical publications including The Lancet, Mayo Clinic Proceedings, and Nature. Tools to study our phase IV findings are available to the public, anonymous and free. On Apr, 08, 2025 141 people who take Amoxicillin and clavulanate potassium and Tylenol together, and have interactions are studied. Amoxicillin and clavulanate potassium has active ingredients of amoxicillin; clavulanate potassium. It is often used in sinusitis. eHealthMe is studying from 6,180 Amoxicillin and clavulanate potassium users for the drug's side effects, drug interactions, effectiveness and more. Check Amoxicillin and clavulanate potassium in the real world. What is Tylenol? Tylenol has active ingredients of acetaminophen. It is often used in pain. eHealthMe is studying from 222,268 Tylenol users for the drug's side effects, drug interactions, effectiveness and more. Check Tylenol in the real world. Female: 0-1 2-9 10-19 20-29 30-39 40-49 50-59 60+ Male: 0-1 2-9 10-19 20-29 30-39 40-49 50-59 60+ Number of Amoxicillin and clavulanate potassium and Tylenol reports submitted per year: What are the common drug interactions of Amoxicillin and clavulanate potassium and Tylenol, by gender? * Click here to view What are the common drug interactions of Amoxicillin and clavulanate potassium and Tylenol, by age (0-1 to 60+)? * Click here to view What are the existing conditions these people have? * Click here to view * Approximation only. Some reports may have incomplete information. Do you take Amoxicillin and clavulanate potassium and Tylenol? - Personalize this study to your gender, age, symptoms and drugs - Predict drug outcomes for up to one year with AI - Get an AI agent to monitor your drugs continuously How the study uses the data? The study uses data from the FDA. It is based on amoxicillin; clavulanate potassium and acetaminophen (the active ingredients of Amoxicillin and clavulanate potassium and Tylenol, respectively), and Amoxicillin and clavulanate potassium and Tylenol (the brand names). Other drugs that have the same active ingredients (e.g. generic drugs) are not considered. Dosage of drugs is not considered in the study. How to use the study? DO NOT STOP MEDICATIONS without first consulting your doctor. If there are any serious or long term adverse effects discovered in the study, discuss the study with your doctor to ensure that proper medication management will be in place if applicable. a b c d e f g h i j k l m n o p q r s t u v w x y z a b c d e f g h i j k l m n o p q r s t u v w x y z a b c d e f g h i j k l m n o p q r s t u v w x y z With medical big data and proven AI/ML algorithms, eHealthMe provides a platform for everyone to run phase IV clinical trials. We study millions of patients and 5,000 more each day. Results of our real-world drug study have been referenced on 800+ peer-reviewed medical publications, including The Lancet, Mayo Clinic Proceedings, and Nature. Our analysis results are available to researchers, health care professionals, patients (testimonials), and software developers (open API). WARNING: Please DO NOT STOP MEDICATIONS without first consulting a physician since doing so could be hazardous to your health. DISCLAIMER: All material available on eHealthMe.com is for informational purposes only, and is not a substitute for medical advice, diagnosis, or treatment provided by a qualified healthcare provider. All information is observation-only. Our phase IV clinical studies alone cannot establish cause-effect relationship. Different individuals may respond to medication in different ways. Every effort has been made to ensure that all information is accurate, up-to-date, and complete, but no guarantee is made to that effect. The use of the eHealthMe site and its content is at your own risk. If you use this eHealthMe study on publication, please acknowledge it with a citation: study title, URL, accessed date. A study for a 79 year old man who takes Olanzapine, Buspar, Mirtazapine - 2 hours ago Drug comparison of Methadone Hydrochloride, Methocarbamol for a 42 year old man - 2 hours ago Drug comparison of Mounjaro, Apple Cider Vinegar for a 63 year old woman - 3 hours ago A study for a 78 year old man who takes Warfarin Sodium, Farxiga - 4 hours ago A study for a 51 year old man who takes Lexapro - 6 hours ago A study for a 61 year old man who takes Methylene Blue, Bupropion Hydrochloride - 6 hours ago A study for a 72 year old man who takes Tamsulosin - 7 hours ago A study for a 70 year old woman who takes Miralax - 7 hours ago A study for a 72 year old man who takes Tamsulosin - 7 hours ago A study for a 61 year old man who takes Bayer Low Dose Aspirin (81mg Aspirin), Escitalopram, Cetrizine Hydrochloride, Alprazolam, Niacin And Niacinamide, Irbesartan, Coq10 - 12 hours ago 8 Topics Found Can you take Tylenol with Amoxicillin Clavulanate as a baby? My 6 month old baby has an ear infection and the dr prescribed a new kind of stronger medication for him today...Clavulin. He just woke up screaming and was a little raspy on his chin & cheeks. We gave him baby Tylenol. What else should we do? ## They can be taken together, with a doctor's approval. The NIH warns that the side effects may include dizziness, headache, non-allergic skin rash, and diarrhea. What are you referring to as raspy? Are you referring to a rash? Has he shown any signs of it being itchy? Can you please post back and clarify? This medication contains Amoxicillin, and Clavulanic Acid, the U.S. name brand for it is Augmentin. NDC code: 43598-004 Has there been any improvement, yet? 1 REPLY Filed under Tylenol Tylenol extra strength and Amox-Clav 875mg Is it safe to take Tylenol extra strength 625mg and Amox-Clav 875 mg together? ## I didn't find any problems or interactions listed between them in the NIH database, as a matter of fact, they are often prescribed together to treat an infection and the pain some of them can cause. The FDA lists the typical side effects as possibly including nausea, dizziness, headache, non-allergic skin rash, and diarrhea. Has your doctor prescribed them both? ## Took my first Amox clav 875mg at 6 pm. My next one is at 6 am the next day. Can I take Tylenol or an Ibuprofen 200 mg in between doses? 2 REPLIES Filed under Tylenol drug interactions with amox tr k clav tylenol I have taken 5 doses so far of amoxicillin/clavulanate (2/day) for a bacterial infection (lymphadenopathy, submandivular -Primary). I have a small crack in a tooth that is now hurting. Can I take Tylenol? How many? Any interactions? ## Hello, Florine! How are you doing? I am sorry that you're in pain. What dosage is the Tylenol? Unless your doctor has instructed otherwise, you should not exceed 1,000mgs every 8 hours, for a maximum of 3,000mgs a day. Taking more risks liver damage, or failure. The FDA lists its typical side effects as possibly including nausea, dizziness, and headache. 1 REPLY Filed under Tylenol Mucinex, Tylenol & Amoxicillin Can I take Mucinex along with Tylenol & Amoxicillin / Clavulanate Potassium Tablets, USP 875 mg/125 mg? ## Hi Nancy, According to the University of Maryland Medical Center (umm.edu / drug interaction tool), there are no interactions listed between Mucinex, Tylenol (acetaminophen) and Amoxicillin. While this doesn't necessarily mean no interactions exist, I personally haven't had any issues taking these together in the past. However, if in doubt, it may be a good idea to double check with your doctor first. I hope this helps! 1 REPLY Filed under Amoxicillin Amox 500mg Clavulan pot I've been taking this for 4 days by IV 1xdaily. 1 more day to go then I start oral for 5 days. Treatment for inflamed kidney. The problem I have is severe headache all the time. Tylenol doesn't help. Starts in my neck behind my left ear it's really tender to touch. Should I maybe talk about an alternative drug? ## Hello, Mel! How are you doing? Has there been any change? That can be a normal side effect of this medication, as reported by the FDA, along with nausea, dizziness, non-allergic skin rash, and diarrhea. These side effects will be more prominent with the stronger IV doses, things should improve once you're taking the lower oral doses, though it may take a week, or so, for things to change. The tenderness behind your ear may be a swollen gland, due to your body fi... 1 REPLY Filed under Tylenol amox tr k clav 600 if you are allergic to cefdinir can you take amox? ## Yes, they are a different class of antibiotics. Cefdinir is a Cephalosporin antibiotic. Amoxicillin with Clavulanic Acid is a generic for Augment, under the Penicillin class of antibiotics. Common side effects may include: nausea, drowsiness, diarrhea and skin rash. Read more: Is there anything else I can help you with? ## I just finished a 7-day cycle of amox-tr-k clav 875-125mg yesterday for a sudden infection in my toe. About 29 hours after I started taking it I got a headache (which I rarely get). It went away w/Tylenol, but returned after Tylenol wore off. I called the doctor the next day & she said it was safe to take Tylenol throughout the cycle, and that if I could stand it, it would be better than starting a new cycle w/a... 2 REPLIES Filed under Cefdinir Relief from pain I began taking amox K clav 875 today for a serious skin
infection. How many days before I can begin feeling some relief from the pain? ## Hello, Linda! How are you doing? I'm sorry that you're in pain. Usually, you should see improvement after taking an antibiotic for a full 3 days. The FDA lists the typical side effects as possibly including nausea, dizziness, headache, non-allergic skin rash and diarrhea. Have you consulted your doctor? It might be okay for you to take Tylenol, or Advil to help, if your doctor approves. 1 REPLY Filed under Tylenol Oval White Pill GGN7 Oval white pill with GGN7 - Is this Amox-Clav 875mg? ## GGN7 is Amoxicillin + Clavulanic Acid 875 mg. This is generic for the Augmentin brand. It is a penicillin antibiotic used to stop bacterial infections. ## LAST TUESDAY AFTERNOON I WENT TO THE DR. AS I HAVE HAD A FEVER FOR OVER A WEEK AND ONLY IBUPROPHEN TOOK THE FEVER DOWN, AND TYLENOL TOOK CARE OF THE PAIN IN MY HIP WHICH IS ARTHRITIS. I HAD AN MRI THAT DECLARED THAT. EVERY NIGHT AND DAY AS I SLEEP I COMPLETELY DRENCH MY PAJAMA SHIRT AND THE PILLOW AND THE BED. I AM NOT IMPROVING. I BOUGHT GATOR AID AS I THOUGHT I WAS LOSING MINERALS. I AM MISERABLE. THE DR. SAID I HAD STREP THROAT ALTHOUGH I HAVE NO PAIN AND SAID I HAD FLUID IN MY EAR AND ALSO THAT WAS NOT IN ANY PAIN. IT HAS BEEN NEARLY A WEEK AND I WONDER IF THIS DRUG SHOULD NO... 4 REPLIES Filed under Amoxicillin Can't find what you're looking for? Certain foods and drinks don't mix well with some medications. A few of these are:Alcohol. Booze can bring on dangerous side effects with many medications, including some drugs for:Before you start a new medication, ask your doctor or pharmacist if it could interact with alcohol. Check the drug label for alcohol warnings, too.Grapefruit or grapefruit juice. Too much of either can affect some drugs like:Grapefruit juice doesn't mix badly with every type of drug in these classes of medications. Check your drug's label or information pamphlet for any warnings about it. Also, ask your doctor or pharmacist if it's safe to have grapefruit or its juice in any amount with your specific drug. If they tell you to stop eating or drinking it, ask if any other fruits or juices might have similar effects on your medicine.Foods with vitamin K, like leafy greens. These can interact with the blood thinner warfarin. You don't have to stop eating them, but it's important to be consistent and not overdo it. Ask your doctor how much food with vitamin K you can have, then eat the same amount around the same times each week.High-potassium foods and drinks, like bananas, salt substitutes, and orange juice. These can affect blood pressure drugs called ACE inhibitors. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also
asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have
any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of
approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal side effects.You also asked about using a saline nasal spray with amoxicillin. That is certainly safe to use as well.Final WordsThanks again for your question! Please reach out again if you have any more questions! Hello, Welcome to clinic.com. A dose of Amoxycillin at 20 years of age is around 1500 mg per day. This total of 1500 mg can be divided into two of approximately 750 mg or three doses of about 500 mg. You have taken the drug recommended in a whole day in six hours, and this will not cause any harm or side effect because
when you take medicine for the first time the drug levels in the body are not maintained fully so if you take the drug in six hours, it will not cause any harm. Acetaminophen (Paracetamol) is a drug for fever, which is usually given with Amoxicillin, and it has no drug reaction with Amoxicillin, so this will also have no effect on your body. In all, I think that you should not worry about this. Do not take any more Amoxicillin today. I hope this will help you.Was this conversation helpful? As a member of the penicillin drug class, amoxicillin is one of the most commonly used antibiotics in the nation. When considering that many bacterial infections are associated with pain and fever, you can probably guess that plenty of circumstances arise when Tylenol and amoxicillin can be helpful. Fortunately, they can be taken together. The problem is that they each have their own side effects and interactions, so it is imperative to read on and understand the details of this drug duo. RELATED: Acetaminophen interactions to avoid | Amoxicillin interactions to avoid To be clear, you can take Tylenol and amoxicillin, which is good news for patients and parents. The drugs don't directly affect each other. This means they don't worsen each other's side effects, and one drug doesn't make the other less effective. For example, suppose you are taking amoxicillin for a sinus infection. In that case, Tylenol is a reasonable option for relieving the associated facial pain or headache; likewise, acetaminophen can be taken as a fever reducer or to reduce body pain from pneumonia treated with amoxicillin. The amoxicillin could cure a urinary tract infection, while Tylenol reduces the associated lower abdominal pain. Can you take Tylenol PM with amoxicillin? Tylenol PM adds diphenhydramine, the active ingredient in brand-name Benadryl, to acetaminophen. Diphenhydramine is an antihistamine that is useful for the treatment of nasal allergic skin rash and diarrhea. Your doctor will track your potassium levels, and they may tell you to cut back on foods with it.St. John's wort. This herbal dietary supplement can affect many medications for heart disease, HIV, depression, and other conditions. It can also affect birth control pills and the cancer drugs irinotecan and imatinib.Always get your doctor's or pharmacist's OK before you try a new supplement. It's also possible for a medication you take to interact with a health condition you have. Some common drug-condition interactions happen between:Antacids and kidney diseaseAntihistamines and certain breathing problems, glaucoma, and enlarged prostateAsthma rescue inhalers and conditions like heart disease, high blood pressure, thyroid disease, and diabetesDecongestants and conditions like heart disease, high blood pressure, thyroid disease, and diabetesSleep aids and certain breathing problems, glaucoma, and enlarged prostate Hello and thanks for reaching out to us!It's not uncommon for someone dealing with an infection to need some symptom relief while they wait for their antibiotic (amoxicillin in this case) to start working.The good news is that it is safe to take either Tylenol (acetaminophen) or Advil (ibuprofen) with amoxicillin as there is no listed or reported interaction between the medications. More DetailAmoxicillin is a "beta-lactam" antibiotic and is structurally similar to penicillin. Unlike penicillin, however, it is considered a "broad spectrum" antibiotic, meaning that it is effective against many types of bacteria (e.g. both gram-positive and gram-negative bacteria).Amoxicillin is a commonly used antibiotic for a variety of infections including sinusitis, bronchitis, and other respiratory tract infections. It is a common need to treat symptoms of pain, headache, and muscle cramps while taking antibiotics such as amoxicillin. Fortunately, most over-the-counter analgesics are safe to take with amoxicillin, including Tylenol (acetaminophen) and Advil (ibuprofen). While there is always a chance of nausea when combining amoxicillin with an analgesic, they are considered safe to use together and should not cause any untoward effects. However, if amoxicillin is causing any stomach-related side effects such as nausea, cramping, or diarrhea, Tylenol may be better tolerated and the preferred option over Advil since NSAIDs, like Advil, tend to cause more gastrointestinal