I'm not a bot

Taking levofloxacin increases the risk that you will develop tendinitis (swelling of a fibrous tissue that connects a bone to a muscle) or have a tendon rupture (tearing of a fibrous tissue that connects a bone to a muscle) during your treatment or for up to several months afterward. These problems may affect tendons in your shoulder, your hand, the back of your ankle, or in other parts of your body. Tendinitis or tendon rupture may happen to people of any age, but the risk is highest in people over 60 years of age. Tell your doctor if you have or have ever had a kidney, heart, or lung transplant; kidney disease; a joint or tendon disorder such as rheumatoid arthritis (a condition in which the body attacks its own joints, causing pain, swelling, and loss of function); or if you participate in regular physical activity. Tell your doctor and pharmacist if you are taking oral or injectable steroids such as dexamethasone, methylprednisolone (Medrol), or prednisone (Rayos). If you experience any of the following symptoms of tendinitis, stop taking levofloxacin, rest, and call your doctor immediately: pain, swelling, tenderness, stiffness, or difficulty in moving a muscle. If you experience any of the following symptoms of tendon rupture, stop taking levofloxacin and get emergency medical treatment: hearing or feeling a snap or pop in a tendon area, bruising after an injury to a tendon area, or inability to move or bear weight on an affected area.Taking levofloxacin may cause changes in sensation and nerve damage that may not go away even after you stop taking levofloxacin. This damage may occur soon after you begin taking levofloxacin. Tell your doctor if you have ever had peripheral neuropathy (a type of nerve damage that causes tingling, numbness, and pain in the hands and feet). If you experience any of the following symptoms, stop taking levofloxacin and call your doctor immediately: numbness, tingling, pain, burning, or weakness in the arms or legs; or a change in your ability to feel light touch, vibrations, pain, heat, or cold. Taking levofloxacin may affect your brain or nervous system and cause serious side effects. This can occur after the first dose of levofloxacin. Tell your doctor if you have or have ever had seizures, epilepsy, cerebral arteriosclerosis (narrowing of blood vessels in or near the brain that can lead to stroke or ministroke), stroke, changed brain structure, or kidney disease. If you experience any of the following symptoms, stop taking levofloxacin and call your doctor immediately: seizures; tremors; dizziness; lightheadedness; headaches that won't go away (with or without blurred vision); difficulty falling asleep or staying asleep; nightmares; not trusting others or feeling that others want to hurt you; hallucinations (seeing things or hearing voices that do not exist); thoughts or actions towards hurting or killing yourself; feeling restless, anxious, nervous, depressed, memory problems, or confused, or other changes in your mood or behavior.Taking levofloxacin may worsen muscle weakness in people with myasthenia gravis (a disorder of the nervous system that causes muscle weakness) and cause severe difficulty breathing or death. Tell your doctor if you have myasthenia gravis. Your doctor may tell you not to take levofloxacin. If you have myasthenia gravis and your doctor tells you that you should take levofloxacin, call your doctor immediately if you experience muscle weakness or difficulty breathing during your treatment.Talk to your doctor about the risks of taking levofloxacin.Your doctor or pharmacist will give you the manufacturer's patient information sheet (Medication Guide) when you begin treatment with levofloxacin. Read the information carefully and ask your doctor or pharmacist if you have any questions. You can also visit the Food and Drug Administration (FDA) website (or the manufacturer's website to obtain the Medication Guide. El levofloxacino es un antibiótico indicado para el tratamiento de infecciones bacterianas, como sinusitis, neumonía, erisipela, forúnculos o infección del tracto urinario, ya que actúa eliminando bacterias del organismo, sensibles a este antibiótico. Este medicamento puede encontrarse en farmacias o droguerías en forma de tabletas de 500 mg o 750 mg, bajo los nombre comerciales Bredelin, Elequine o Sibofix, por ejemplo, o como genérico “levofloxacino”, y debe usarse bajo indicación médica, siendo vendido solo con la presentación de receta médica. Además, el levofloxacino también puede encontrarse en presentación inyectable, la cual debe ser aplicada por un enfermero en hospitales. Para qué sirve El levofloxacino está indicado para el tratamiento de infecciones bacterianas como: Sinusitis aguda; Neumonía; Impétigo; Abscesos en la piel; Forúnculo; Celulitis bacteriana; Erisipela; Osteomielitis; Infecciones del tracto urinario, como pielonefritis; Infecciones intraabdominales; Infección generalizada o sepsis. Además, el levofloxacino puede indicarse para el tratamiento de exacerbaciones agudas de bronquitis crónica. ¿El levofloxacino puede usarse para COVID-19? El levofloxacino puede ser indicado por el médico en personas con COVID-19 y que tengan al mismo tiempo neumonía bacteriana, ya que este medicamento ayuda a combatir bacterias. Sin embargo, el levofloxacino no tiene ninguna acción contra virus, incluso el coronavirus, no estando indicado cuando no hay infecciones bacterianas asociadas. Vea los principales medicamentos indicados para el tratamiento de COVID-19. Cómo tomar levofloxacino El levofloxacino en presentación de tabletas debe tomarse por vía oral, con un vaso de agua, antes o después de una comida, en los horarios establecidos por el médico. Sin embargo, personas que utilizan medicamentos como sales de hierro, antiácidos o sucralfato, deben tomar la tableta de levofloxacino 2 horas antes o 2 horas después del uso con este tipo de medicamentos. La dosis de levofloxacino varía de acuerdo al tipo de infección a tratar, siendo que las dosis normalmente recomendadas para adultos, son: El tiempo de tratamiento con levofloxacino debe ser siempre orientado por el médico de acuerdo a la condición a tratar. Posibles efectos secundarios Los efectos secundarios más comunes que pueden surgir durante el tratamiento con levofloxacino son náuseas, vómitos, diarrea, dolor de cabeza, mareos, insomnio o dolor abdominal. Además, también puede haber ruptura del tendón de Aquiles, convulsiones, alucinaciones, agitación, ansiedad, alteración en el gusto, pancreatitis, formación de bolitas en la piel, comezón, urticaria, dolor en el cuerpo, debilidad muscular, malestar, fiebre o alteraciones en la función del hígado. Contraindicaciones El levofloxacino debe evitarse en niños, adolescentes, mujeres embarazadas o que estén lactando, así como en personas que tengan epilepsia, antecedentes de tendinitis o ruptura en el tendón. Además, este medicamento no debe usarse en personas que tengan alergia al levofloxacino o a otras fluoroquinolonas, como ciprofloxacino o norfloxacino, por ejemplo, o a cualquier otro componente de la fórmula. New search Hide text from GuidelinesJ ANTIINFECTIVES FOR SYSTEMIC USE Antiinfectives are also classified in the following groups:A01AB Antiinfectives and antiseptics for local oral treatmentA02BD Combinations for eradication of Helicobacter pyloriA07A Intestinal antiinfectivesD01 Antifungals for dermatological useD06 Antibiotics and chemotherapeutics for dermatological useD07C Corticosteroids, combinations with antibioticsD09AA Ointment dressings with antiinfectivesD10AF Antiinfectives for treatment of acneG01 Gynecological antiinfectives and antisepticsP Antiparasitic products, insecticides and repellentsR02AB AntibioticsS01/S02/S03 Eye and ear preparations with antiinfectivesEven systemically administered antibacterials and antimycotics may be classified in other groups if their target is exclusively local, e.g. the skin - D01 - Antifungals for dermatological use.Inhaled antiinfectives are classified in J.The DDDs for the antiinfectives are as a main rule based on the use in infections of moderate severity. However, some antiinfectives are only used in severe infections and their DDDs are assigned accordingly. The DDDs assigned are based on daily treatment. The duration of the treatment periods is not taken into consideration. For antiinfectives given in a high initially starting dose followed by a lower daily "maintenance" dose, the DDDs are based on the "maintenance" dose if the total duration of the treat-ment course is more than one week. If, however, the treatment course is 7 days or less, the DDDs are assigned according to the average daily dose i.e. the total course dose divided by the number of treatment days (e.g azithromycin). J01 ANTIBACTERIALS FOR SYSTEMIC USE This group comprises antibacterials for systemic use, except antimycobacterials, which are classified in J04. The antibacterials are classified according to their mode of action and chemistry.Combinations of two or more systemic antibacterials from different third levels are classified in J01R, except combinations of sulfonamides and trimethoprim, which are classified at a separate 4th level, J01EE.Combinations of antibacterials and tuberculostatics are classified in J04AM.Combinations of antibacterials with antiparasitic agents are classified in J01R.Combinations of antibacterials with other drugs, including local anesthetics or vitamins, are classified at separate 5th levels in the respective antibacterial group by using the 50-series.Inhaled antiinfectives are classified here based on the fact that preparations for inhalation can not be separated from preparations for injection. J01R COMBINATIONS OF ANTIBACTERIALS This group comprises combinations of two or more antibacterials for systemic use from different ATC 3rd levels.The detailed classification of antibacterial combinations in J01RA is based on the general concern with the use of antibacterials worldwide and the need for drug monitoring, incl. mapping of the use with resistance patterns.Combinations of antibacterials with antiparasitic agents are classified here. J01RA Combinations of antibacterials Combinations of urinary antiseptics and antiinfectives are classified in J01RA02. ATC code Name DDD U Adm.R Note J01RA05 levofloxacin and ornidazole List of abbreviations Share — copy and redistribute the material in any medium or format for any purpose, even commercially. Adapt — remix, transform, and build upon the material for any purpose, even commercially. The licensor cannot revoke these freedoms as long as you follow the license terms. Attribution — You must give appropriate credit , provide a link to the license, and indicate if changes were made . You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use. ShareAlike — If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original. No additional restrictions — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits. You do not have to comply with the license for elements of the material in the public domain or where your use is permitted by an applicable exception or limitation . No warranties are given. The license may not give you all of the permissions necessary for your intended use. For example, other rights such as publicity, privacy, or moral rights may limit how you use the material. El levofloxacino es un antibiótico indicado para el tratamiento de infecciones bacterianas, como sinusitis, neumonía, erisipela, forúnculos o infección del tracto urinario, ya que actúa eliminando bacterias del organismo, sensibles a este antibiótico. Este medicamento puede encontrarse en farmacias o droguerías en forma de tabletas de 500 mg o 750 mg, bajo los nombre comerciales Bredelin, Elequine o Sibofix, por ejemplo, o como genérico “levofloxacino”, y debe usarse bajo indicación médica, siendo vendido solo con la presentación de receta médica. Además, el levofloxacino también puede encontrarse en presentación inyectable, la cual debe ser aplicada por un enfermero en hospitales. Para qué sirve El levofloxacino está indicado para el tratamiento de infecciones bacterianas como: Sinusitis aguda; Neumonía; Impétigo; Abscesos en la piel; Forúnculo; Celulitis bacteriana; Erisipela; Osteomielitis; Infecciones del tracto urinario, como pielonefritis; Infecciones intraabdominales; Infección generalizada o sepsis. Además, el levofloxacino puede indicarse para el tratamiento de exacerbaciones agudas de bronquitis crónica. ¿El levofloxacino puede usarse para COVID-19? El levofloxacino puede ser indicado por el médico en personas con COVID-19 y que tengan al mismo tiempo neumonía bacteriana, ya que este medicamento ayuda a combatir bacterias. Sin embargo, el levofloxacino no tiene ninguna acción contra virus, incluso el coronavirus, no estando indicado cuando no hay infecciones bacterianas asociadas. Vea los principales medicamentos indicados para el tratamiento de COVID-19. Cómo tomar levofloxacino El levofloxacino en presentación de tabletas debe tomarse por vía oral, con un vaso de agua, antes o después de una comida, en los horarios establecidos por el médico. Sin embargo, personas que utilizan medicamentos como sales de hierro, antiácidos o sucralfato, deben tomar la tableta de levofloxacino 2 horas antes o 2 horas después del uso con este tipo de medicamentos. La dosis de levofloxacino varía de acuerdo al tipo de infección a tratar, siendo que las dosis normalmente recomendadas para adultos, son: El tiempo de tratamiento con levofloxacino debe ser siempre orientado por el médico de acuerdo a la condición a tratar. Posibles efectos secundarios Los efectos secundarios más comunes que pueden surgir durante el tratamiento con levofloxacino son náuseas, vómitos, diarrea, dolor de cabeza, mareos, insomnio o dolor abdominal. Además, también puede haber ruptura del tendón de Aquiles, convulsiones, alucinaciones, agitación, ansiedad, alteración en el gusto, pancreatitis, formación de bolitas en la piel, comezón, urticaria, dolor en el cuerpo, debilidad muscular, malestar, fiebre o alteraciones en la función del hígado. Contraindicaciones El levofloxacino debe evitarse en niños, adolescentes, mujeres embarazadas o que estén lactando, así como en personas que tengan epilepsia, antecedentes de tendinitis o ruptura en el tendón. Además, este medicamento no debe usarse en personas que tengan alergia al levofloxacino o a otras fluoroquinolonas, como ciprofloxacino o norfloxacino, por ejemplo, o a cualquier otro componente de la fórmula